

Distributed by:



A division of Pe-Bee Agri (Pty) Ltd
cnr R23 Benoni Road & R550 Nigel Road
Brakpan District.

Telephone: 011 812 9800
www.protek.co.za

Risk and Safety Information.

To avoid risk to human health and the environment, comply with the instructions for use.

GHS Signal word: **WARNING**
HAZARD STATEMENT:

H303: May be harmful if swallowed.
H313: May be harmful in contact with skin.
H332: May be harmful if inhaled.

DISCLAIMER: Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONARY STATEMENT:

P101: If medical advice is needed, have product container or label at hand.
P102: Keep out of reach of children.

P103: Read carefully and follow instructions.

PREVENTION STATEMENT:

P270: Do not eat, drink, or smoke when using this product.
P280: Wear protective gloves.

WARNINGS

HANDLE WITH EXTREME CARE. Prevent access to bait by children and animals. Store away from food and feed. Poisonous if swallowed and on prolonged contact with skin. Toxic to fish and wildlife. In case of poisoning, call a doctor or a veterinarian and make this label available to him.

PRECAUTIONS

Avoid contamination of feed, food, eating utensils and water. Set bait stations where they will be inaccessible to children, pets, livestock, and other non-target animals. Wash hands and exposed skin after handling the poison. Do not eat, drink or smoke while handling the poison. Destroy empty container and do not use for any other purpose. Bury dead rodents and any uneaten bait at the end of treatment.

SCIENTIFIC™



SUPA-KILL

RAT & MOUSE POISON BAIT

READY-TO-USE, GRAIN BAIT.

Reg. No. SA: L5198 Act 36/1947 Namibia: N-AR 1144

READ THE LABEL CAREFULLY AND FOLLOW ALL INSTRUCTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

CONTROLS: ROOF RATS, NORWAY RATS AND COMMON HOUSE MICE.

USUALLY AFTER A SINGLE FEED.

EFFECTIVE AGAINST WARFARIN RATS AND MICE.

If medical advice is needed, have product container or label at hand.

Active Ingredient: BROMADIOLONE: 0.05 g/kg.



WARNING



HAZARD STATEMENT:

May be harmful if swallowed.

May be harmful in contact with skin.

PRECAUTIONARY STATEMENT:

Keep out of reach of children.

Read carefully and follow instructions.

Registered and Manufactured by:

SCIENTIFIC SUPA-KILL c.c.

CK 1991/27877/23

P.O.Box 8020, Edleen 1625,

Tel. No. (011) 394-3941/2

www.supa-kill.com

EMERGENCY POISONING

INFORMATION:


Tygerberg Poison Information Centre:

0861555777

3kg

UN No.: 3027

Batch No:



Date of Manufacture:

SYMPTOMS OF HUMAN POISONING

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around the elbow and knee joints. Blood in the urine and faeces.

FIRST AID TREATMENT

If swallowed, call the Tygerberg Poison Information Centre: 0861555777 for treatment advice. Only induce vomiting if advised to do so or if it is less than three hours since the bait was eaten. Take the person to a hospital or a doctor IMMEDIATELY and show the product label. In the case of animal poisonings consult a veterinarian.

NOTE TO PHYSICIAN OR VETERINARIAN:

Bromadiolone is a delayed-action anticoagulant. Perform stomach wash if a large amount of bait was ingested and less than 4 hours have elapsed. Determine prothrombin time 12 to 18 hours after ingestion and if elevated, give Vitamin K1 intravenously in divided doses. In severe cases transfusions of fresh blood may be necessary. CONTACT A POISON INFORMATION CENTRE.

Relevant Compounds

Compound	Weight %
Bromadiolone	0.005
Bitrex	0.001

DIRECTIONS FOR USE

For use in the home, on the farm and industrial premises. Supa-Kill Rat and Mouse poison can kill rodents after a single feed. Death will take place within 4 to 12 days. **EFFECTIVE AGAINST WARFARIN RESISTANT RATS AND MICE. ALWAYS PLACE SUPA-KILL RAT & MOUSE POISON IN A COVERED BAIT STATION TO PREVENT ACCESS TO BAIT BY CHILDREN AND ANIMALS.** Bait points should be inspected and replenished every 7 days until feeding ceases.

USE ONLY AS DIRECTED

NORWAY AND ROOF RATS

Place 50-75g Supa-Kill Rat & Mouse poison bait in Supa-Kill bait stations, along walls and active rat runs and in sheltered areas where rats are known to live or feed.

MICE

Place 20g Supa-Kill Rat & Mouse poison bait in bait stations every 2 to 4 meters along walls or runways, in nests and in sheltered areas where mice are known to live or feed. The single feeding action of Supa-Kill Rat and Mouse poison is particularly effective against mice with their small appetites and sporadic feeding behaviour.



6 009705 422289